



# HEALTHY HEART

VOLUME-14 | ISSUE-169 | DECEMBER 05, 2023

Price : ₹ 5/-

Honorary Editor:

**Dr. Hemang Baxi**

Interventional Cardiologist



**From the Desk of Hon. Editor:**

Dear Friends,

It is a great pleasure to interact with you periodically and share scientific knowledge.

In this issue, I have tried to summarise Complications of Acute Myocardial Infarction (AMI).

Although the incidence of various post MI complications have decreased significantly with the advent of newer pharmacological agents, early reperfusion techniques (Pharmacological or mechanical), prognosis remains dismal for several important

post-AMI complications.

We will be discussing various clinical aspects & management issues of all these complications.

**Corresponding Email : [hemang.baxi@cims.me](mailto:hemang.baxi@cims.me)**

## Complications of Acute Myocardial Infarction

### Introduction

Each year, approximately ten million The patients in India suffer an AMI, leading directly to 8,00,000 deaths. Despite new pharmacologic agents, the emergence of reperfusion therapy, and the growing acceptance of early revascularization, prognosis remains dismal for several important post-AMI complications. Thus, to achieve the best possible outcomes, cardiology providers need to be able to recognize, diagnose, and treat AMI complication, including : 1) cardiogenic shock, 2) right ventricular (RV) infarction/ischemia, 3) ischemic mitral valve regurgitation (MR), 4) ventricular septal defect (VSD), and 5) left ventricular (LV) free wall rupture.

### Cardiogenic shock

#### Incidence

Cardiogenic shock remains the leading cause of death for patient hospitalized with AMI, and complicates 6-7 % of acute

ST-elevation and non-ST-elevation MIs (NSTEMIs). In the Global Utilization of Streptokinase and Tissue Plasminogen Activator for Occluded Coronary Arteries (GUSTO) IIb trial, 4.2 % of ST-segment elevation patients developed cardiogenic shock compared to only 2.5 % of patients with non-ST-segment elevation; however, mortality secondary to cardiogenic shock is similarly high among patients with and without ST-segment elevation.

Severe dysfunction of the LV due to infarction and ischemia is the most common cause of cardiogenic shock. However, it can also be precipitated by other complications of AMI such as RV dysfunction, acute MR, VSD, or cardiac free wall rupture.

#### Clinical presentation

In addition to the MI symptoms, patients with cardiogenic shock will develop respiratory distress and signs of

#### Cardiologists

Dr Vipul Kapoor (M) +91-98240 99848	Dr Urnil Shah (M) +91-98250 66939
Dr Tejas V. Patel (M) +91-89403 05130	Dr Hemang Baxi (M) +91-98250 30111
Dr Hiren Kevadiya (M) +91-98254 65205	Dr Anish Chandarana (M) +91-98250 96922
Dr Gunvant Patel (M) +91-98240 61266	Dr Ajay Naik (M) +91-98250 82666
Dr Keyur Parikh (M) +91-98250 26999	Dr Satya Gupta (M) +91-99250 45780
Dr Milan Chag (M) +91-98240 22107	Dr Vineet Sankhla (M) +91-99250 15056

#### Congenital & Structural Heart Disease Specialist

Dr Kashyap Sheth (M) +91-99246 12288	Dr Milan Chag (M) +91-98240 22107
--------------------------------------	-----------------------------------

#### Cardiothoracic & Vascular Surgeons

Dr Dhiren Shah (M) +91-98255 75933	Dr Dhaval Naik (M) +91-90991 11133
Dr Amit Chandan (M) +91-96990 84097	Dr Kishore Gupta (M) +91-99142 81008
Dr Nikunj Vyas (M) +91-73531 65955	

#### Paediatric & Structural Heart Surgeons

Dr Shaunak Shah (M) +91-98250 44502
-------------------------------------

#### Cardiovascular, Thoracic & Thoracoscopic Surgeon

Dr. Pranav Modi (M) +91-99240 84700
-------------------------------------

#### Cardiac Anaesthetists

Dr. Niren Bhavsar (M) +91-98795 71917	Dr. Hiren Dholakia (M) +91-95863 75818
Dr. Chintan Sheth (M) +91-91732 04454	

#### Cardiac Electrophysiologist

Dr. Ajay Naik (M) +91-98250 82666	Dr. Hiren Kevadiya (M) +91-98254 65205
-----------------------------------	--



hypoperfusion including cool, clammy extremities, oliguria, and mental status changes from cerebral hypoperfusion. Physical exam reveals hypotension, rales, S3 gallop, and cold extremities. Hemodynamically, the systemic systolic pressure is lower than 90 mm Hg, the cardiac index is lower than 1.8 L/min/m<sup>2</sup>, and systemic vascular resistance and pulmonary artery wedge pressure are elevated. Laboratory data reveal lactic acidosis, elevated blood urea nitrogen/creatinine, and arterial hypoxemia. Chest X-ray reveals pulmonary edema.

## Diagnosis

In the Should we emergently revascularize occluded coronaries for cardiogenic shock (SHOCK) trial, the diagnosis of cardiogenic shock was confirmed by both clinical and hemodynamic criteria. The clinical criteria were:

- Hypotension (systolic blood pressure <90 mm Hg for at least 30 minutes)
- End-organ hypoperfusion (cool extremities or a urine output < 30 mL/h, and a heart rate > 60 beats/min)

## The hemodynamic criteria were :

- Cardiac index < 2.2 L/min per square meter of body-surface area
- Pulmonary-capillary wedge pressure of at least 15 mm Hg

The median time from MI to onset of cardiogenic shock in the SHOCK trial was 5.5 hours, with 75 % of patients developing shock within 24 hours.

## Management

The goals of the therapy for cardiogenic shock included hemodynamic stabilization to ensure tissue perfusion and prompt assessment for reversible cause of cardiogenic shock. Intra-aortic balloon pump (IABP) counterpulsation should be considered. Positives inotropes (dobutamine and dopamine) should be added if necessary to improve cardiac output and hemodynamics. Diuretics should be used to treat pulmonary congestion and enhance oxygenation.

Reperfusion therapy is likely to improve post-AMI cardiogenic shock mortality. This has been suggested by GUSTO-1 trial subgroup analysis, which studied 1,320 patients with shock. Reperfusion therapy was associated with reduced in-hospital (50 %) and one year mortality. The SHOCK trial is the largest and most important randomized trial evaluating early revascularization in patients with cardiogenic shock. Patients with shock due to LV failure complicating MI were randomized to emergency revascularization or initial medical stabilization.

Revascularization was accomplished by either coronary artery bypass grafting (CABG) or angioplasty. The primary endpoint was mortality from all causes at 30 days, and the secondary endpoint was six-month survival. Revascularization was associated with a nonsignificant mortality reduction at 30 days (absolute risk reduction 9 % relative risk reduction 17 %); however, the six-month mortality rate was

significantly reduced (50.3 % vs 63.1 %,  $P = 0.027$ ). Subgroup analysis suggested that patients younger than age 75 benefited proportionally more from early reperfusion strategy.

## RV infarction /ischemia

### Incidence

Approximately 50 % of patients who develop an inferior MI will also experience a RV infarction/ischemia, although only about 10 % of those




Figure-1 : The downward spiral in cardiogenic shock

infarcts will be clinically and hemodynamically significant.

### Clinical Presentation

The clinical triad of hypotension, clear lung fields, and elevated jugular venous pressure (JVP) in a patient with an inferior infarction is virtually pathognomonic for RV infarction / ischemia. Although quite specific, this triad has a sensitivity of <25%. Physical exam reveals elevated JVP and RV S3



with clear lungs (in patients without concomitant LV dysfunction). Kussmaul's sign (inspiratory increase in JVP) can also be seen. The predominant atrial descent is systolic (x descent) and the diastolic (y) descent is blunted, reflecting an increased resistance to diastolic filling.

## Diagnosis

**ECG:** Lead V4R is the single best lead for diagnosis, 1 mm ST elevation in lead V4R is approximately 70% sensitive and 100% specific for acute RV infarction. Thus, right-sided ECGs should be done in all patients with acute inferior infarction.

**Echocardiography** is the study of choice for RV infarction. It provides information about the size and function of the RV.

**Pulmonary artery catheterization** reveals elevated RA and RV pressure and low pulmonary capillary wedge pressure (unless there is concomitant LV failure).

## Management

Optimal management of RV involvement in AMI or cardiogenic shock requires early recognition. Once suspected, the treatment should be directed at early maintenance of RV preload, reduction of RV afterload, inotropic support of the dysfunctional RV, and early reperfusion. Intravenous fluids (usually normal saline) should be given to enhance the RV and eventually the LV filling pressure. Agents that decrease the RV preload such as nitrates, diuretics and morphine may worsen hemodynamic condition and should be restricted. Inotropes can be used in cases where fluid resuscitation is not sufficient to improve the hemodynamic status of the patient.

Maintenance of atrioventricular synchrony is also important to optimize RV filling. Most importantly, patients with RV involvement in either LV or RV shock can benefit as much from early revascularization as other patients with cardiogenic shock. Patients who undergo successful reperfusion have improved RV function reduced 30-day mortality.

## Acute Mitral Valve Regurgitation

MR following AMI results from regional or global LV ischemia, papillary muscle dysfunction or papillary muscle rupture. Mitral regurgitation is fairly common following AMI and will persist in 15% cases. Papillary muscle rupture causing MR occurs much less frequently and complicates approximately 2 % of AMIs, often leading to cardiogenic shock and worse outcome. In the SHOCK trial registry, acute severe MR was the cause of shock.

## Clinical manifestations

The diagnosis of ischemia-induced acute MR and papillary muscle rupture is often difficult. Severe MR and heart failure symptoms may develop following a small or moderate-sized AMI. Most patients develop acute hypotension and flash pulmonary edema with a new holosystolic murmur. Although the murmur is often loud, a palpable thrill is generally not present (in contrast to VSD). The intensity of the murmur frequently does not correlate with the severity of the regurgitant flow. If the LV function is severely impaired or if the left atrial pressure is very high, the murmur may be of low intensity or even absent. If a pulmonary artery catheter is already in place, the pulmonary capillary pressure

tracing usually shows giant V waves secondary to decreased left atrial compliance in the setting of acute regurgitation. This phenomenon is also seen in acute VSD.

## Diagnosis

The diagnosis is usually suggested by the combination of hemodynamic compromise and a new systolic murmur in the setting of an AMI. The main differential diagnosis is postinfarction VSD.

Echocardiography with color flow Doppler is valuable to confirm the diagnosis and mechanism of ischemia-induced acute MR.

## Treatment

Emergent arterial vasodilators such as nitroprusside may improve the hemodynamic status temporarily because it decreases systemic vascular resistance. This reduces the regurgitant fraction and increases the forward stroke volume and cardiac output. Most cases of ischemia-induced MR resolve or improve without surgery following reperfusion/revascularization and treatment of the underlying ischemia/infarction. Surgery is usually required in presence of papillary muscle rupture, much less commonly in the absence of rupture. Emergent surgical intervention remains the treatment of choice for papillary muscle rupture. Observational data suggest that surgery for acute MR should be performed acutely, even in patients who appear to stabilize with medical therapy, because its unpredictable. The perioperative mortality associated with postinfarct





papillary muscle rupture is high (27%), but two-thirds of the survivors are alive at seven years. The only factor that improved both immediate and long-term

survival was the concomitant performance of CABG.

## Rupture of the interventricular septum (Post AMI VSD)

### Incidence

Rupture of the interventricular septum occurred in about 2 % of all acute infarctions in the prethrombolytic era. This figure has decreased to 0.2 % in patients treated with reperfusion therapy. It may develop as early as 24 hours after MI, but typically occurs 3-5 days after the acute event. In the GUSTO-I trial, VSD was associated with a very high 30-day mortality (74 % vs. 7 % for patients with and without VSD)

### Clinical manifestations

Patients with a ruptured septum usually present with hypotension, congestive heart failure (CHF), and a new murmur. The murmur is harsh, loud, and holosystolic, best heard at the lower left sternal borders and accompanied by a palpable thrill in 50% of cases. VSD can present as cardiogenic shock.

### Diagnosis

Two-dimensional TTE with color flow imaging is the recommended test to diagnose post-AMI VSD.

### Management

The key to the management of postinfarction VSD is early recognition and early surgical intervention. Unless

there is significant aortic regurgitation, an IABP should be inserted emergently as bridge to a surgical procedure. Early surgical closure is recommended and




Figure-2 : Posterior ventricular septal rupture (VSD)

likely yields better results than attempting to wait for days, weeks until the conditions for surgery improve. In the SHOCK trial registry, surgical repair was performed on 31 patients with rupture, of whom six (19 %) survived. Of the 24 managed medically, only one survived. In the GUSTO-I trial, surgical repair was performed in 40 percent of patients with a VSD within a median of 3.5 days after onset of infarction symptoms. The 30-day and one-year survival for those undergoing surgery was 53 % and 47 %, respectively, while it was only 6 % and 3 % for those treated medically.

In summary, postinfarct VSD carries a very high mortality, and the key to the management of these patients is rapid stabilization and prompt surgery.

## Rupture of the LV free wall

### Incidence

Rupture of the free ventricular wall is almost always fatal. Fortunately, reperfusion therapy has decreased the incidence of free wall rupture (0.2 %).

More than one-half of myocardial rupture occurs within the first five days after MI. Independent predictors of rupture were anterior location of the infarction, age >70, and female sex.

### Clinical presentation

The most common presentation of myocardial rupture is cardiac arrest leading to sudden death. Electro-mechanical dissociation is usually the underlying cardiac rhythm. Premonitory chest discomfort and intermittent bradycardia signal impending rupture in most cases. Complete rupture of the LV free wall usually leads to hemopericardium and death from acute cardiac tamponade.

### Diagnosis

High clinical suspicion and immediate echocardiography during resuscitation with demonstration of hemopericardium and cardiac tamponade is essential for diagnosis.

### Treatment

Survival of AMI patients with LV free wall rupture and cardiogenic shock is dismal. Very few patients can be resuscitated or temporarily stabilized in order to be considered for surgical repair. Aggressive volume replacement with blood transfusion and intravenous fluids, pressor agents, and emergent pericardiocentesis may be considered if LV rupture is suspected or diagnosed. Still, medical therapy has no role in the treatment for acute LV wall rupture. The only hope for these patients is emergency surgical repair, thus emphasizing the importance of early recognition, diagnosis, and surgical triage.



[cims.org](https://cims.org)



# INSTITUTE OF CARDIAC SCIENCES

One-stop solution for any cardiac disease

**PIONEER IN GUJARAT**

- Primary Angioplasty, IVUS, OCT, IVL
- Neonatal & Pediatric Intervention and Surgery
- EP & RF - Ablation Therapy
- Transcatheter Aortic Valve Implantation (TAVI)



- Heart Failure & Artificial Heart Program
- Minimally Invasive Cardiac Surgery (MICS)
- Heart and Lung Transplant
- Valvular and Structural Heart Disease Therapy



## ONE OF INDIA'S LARGEST TEAM OF HEART CARE EXPERTS

**TOP ROW LEFT TO RIGHT:** Dr Vipul Ahir | Dhanyata Dholakia | Dr Chintan Sheth | Dr Niren Bhavsar  
Dr Nikunj Vyas | Dr Shaunak Shah | Dr Dhiren Shah | Dr Dhaval Naik | Dr Amit Chandan | Dr Pranav Modi  
Dr Kishore Gupta | Dr Hiren Dholakia | Ulhas Padiyar | Akash Rajawat | Dr Guntant Patel

**BOTTOM ROW LEFT TO RIGHT:** Dr Tejas V. Patel | Dr Satya Gupta | Dr Urmil Shah | Dr Anish Chandarana  
Dr Keyur Parikh | Dr Milan Chag | Dr Ajay Naik | Dr Hemang Baxi | Dr Hiren Kevadiya | Dr Vipul Kapoor | Dr Kashyap Sheth



**American College of Cardiology (ACC)**  
Centre of Excellence  
**ONE OF THE FIRST IN INDIA**



**Marengo CIMS Hospital**

Off. Science City Road, Sola, Ahmedabad - 380060  
Email: [marengocims.info@marengoaisa.com](mailto:marengocims.info@marengoaisa.com)



For emergency or appointment,  
**1800 309 9999**



# JIC 2024

Joint International Conference

**JANUARY 05 - 07**

20<sup>th</sup> Annual Scientific Symposium

29<sup>th</sup> Year of Academics

## JIC 2024 : PROGRAM HIGHLIGHTS



### DAY 1 JANUARY 5, 2024

- **Internal Medicine Symposium**
- **Satellite Session**
  - An Evening of Pharmacology & Therapeutics
  - 10 Points to Remember for Physicians



### DAY 2 JANUARY 6, 2024

- **Cardiovascular Symposium**
- **Critical Care (Airway Workshop)**
- **Satellite Session**
  - Gastrology Update
  - Neurology (Stroke) Update
  - Infectious Diseases
  - Back to Basics



### DAY 3 JANUARY 7, 2024

- **Cardiovascular Symposium**
- **Critical Care / Pulmonology Symposium**
- **Gynecology And Infertility Update**
- **Nephrology Symposium**
- **JIC Student Quiz**

Register for JIC 2024  
And Get Informative  
and worth while Satellite  
Sessions Absolutely

**FREE**



**JIC 2024**  
Joint International Conference  
**JANUARY 05 - 07**

**20<sup>th</sup> Annual Scientific Symposium**  
**29<sup>th</sup> Year of Academics**

## REGISTRATION FORM - JIC 2024

PLEASE NOTE THAT IT IS MANDATORY TO PROVIDE ALL THE INFORMATION.

PLEASE FILL IN CAPITAL LETTERS

**FOR MD PHYSICIANS**  
**₹ 2,000/-** \*  
Only

**FOR MD STUDENTS**  
**₹ 1,000/-** \*  
Only

Full Name : \_\_\_\_\_  
Qualification : \_\_\_\_\_  
Speciality : \_\_\_\_\_  
Resi. Address : \_\_\_\_\_  
City : \_\_\_\_\_ Pin Code : \_\_\_\_\_  
Mobile No. : \_\_\_\_\_  
Email : \_\_\_\_\_

Cheque or DD's to be made A/C payee and in the name of  
'CIMS Hospital Pvt. Ltd.' Kindly mail the cheque/DD to our office.

**Note : Hotel Accommodation is optional for you & your Spouse.**  
**Two Night hotel accommodation charges - for you Rs. 5000/- |**  
**and for Spouse Rs. 5000/-**  
**(Please Send Separate Cheque)**

### PAYMENT DETAILS

₹ : \_\_\_\_\_ in words :

DD/Cheque : \_\_\_\_\_ Date : \_\_\_\_ / \_\_\_\_ / \_\_\_\_

### FOR REGISTRATION INFORMATION

(M) +91-90990 66538  
(M) +91-98251 08257

Organized by

GMERS  
Medical College,  
Sola,  
Ahmedabad

Supported by



CIMSRE

Care Institute  
Medical Society  
for Research  
and Education

In Association with



Conference Secretariat

Marengo CIMS Hospital, Off. Science City Road, Sola, Ahmedabad -380060  
(M) +91-90990 66538 | +91-98251 08257  
Email : marengocims.communication@marengoasia.com | www.jicindia.org





## HEALTHY HEART

VOLUME-14 | ISSUE-169 | DECEMBER 05, 2023

Healthy Heart Registered under **RNI No. GUJENG/2008/28043**

Published on 5<sup>th</sup> of every month

Permitted to post at PSO, Ahmedabad-380002 on the 12<sup>th</sup> to 17<sup>th</sup> of every month under

Postal Registration No. **GAMC-1725/2021-23** issued by SSP Ahmedabad valid upto 31<sup>st</sup> December, 2023

Licence to Post Without Prepayment No. **PMG/NG/055/2021-23** valid upto 31<sup>st</sup> December, 2023

### If undelivered Please Return to :

CIMS Hospital,  
Nr. Shukan Mall, Off Science City Road,  
Sola, Ahmedabad-380060.  
Call : 1800 309 9999

**Subscribe "Healthy Heart"** : Get your "Healthy Heart", the information of the latest medical updates only ` 60/- for one year.

To subscribe pay ` 60/- in cash or cheque/DD at CIMS Hospital Pvt. Ltd. Nr. Shukan Mall, Off Science City Road, Sola, Ahmedabad-380060. Phone : +91-79-4805 2823. Cheque/DD should be in the name of : **"CIMS Hospital Pvt. Ltd."**

Please provide your **complete postal address with pincode, phone, mobile and email id** along with your subscription

**SCAN TO  
REGISTER NOW**



**JIC 2024**  
Joint International Conference

**JANUARY 05 - 07**

20<sup>th</sup> Annual Scientific Symposium  
29<sup>th</sup> Year of Academics

**REGISTRATIONS ARE OPEN  
FOR IN-AUDITORIUM MEET WITH LIMITED SEATS.**

**Register during these 3 days for JIC 2024.**

**THE PRIME LEARNING DESTINATION - JOIN US**



**Advanced Multi-Specialty Symposiums**



**Case Based Discussion By Experts**



**Interactive lectures and Debates**

**For MD Physicians**

**₹ 2,000/-\* Only**

**For MD Students**

**₹ 1,000/-\* Only**

CIMS Hospital : Regd Office: Plot No.67/1, Opp. Panchamrut Bungalows, Nr. Shukan Mall, Off Science City Road, Sola, Ahmedabad - 380060.

Call : 1800 309 9999

CIMS Hospital Pvt. Ltd. | CIN : U85110GJ2001PTC039962 | [marengocims.info@marengoasia.com](mailto:marengocims.info@marengoasia.com) | [www.cims.org](http://www.cims.org)

Printed, Published and Edited by Dr. Keyur Parikh on behalf of the CIMS Hospital

Printed at Hari Om Printery, 15/1, Nagori Estate, Opp. E.S.I. Dispensary, Dudheshwar Road, Ahmedabad-380004.

Published from CIMS Hospital, Nr. Shukan Mall, Off Science City Road, Sola, Ahmedabad-380060.