


# Healthy Heart

Volume-3 | Issue-30 | May 5, 2012



Price : ₹ 5/-

**Honorary Editor :**  
Dr. Hemang Baxi

#### Cardiologists

Dr. Anish Chandarana  
(M) +91-98250 96922  
Dr. Ajay Naik  
(M) +91-98250 82666  
Dr. Satya Gupta  
(M) +91-99250 45780  
Dr. Joyal Shah  
(M) +91-98253 19645  
Dr. Ravi Singhvie  
(M) +91-98251 43975  
Dr. Gunvant Patel  
(M) +91-98240 61266  
Dr. Keyur Parikh  
(M) +91-98250 26999  
Dr. Milan Chag  
(M) +91-98240 22107  
Dr. Urmil Shah  
(M) +91-98250 66939  
Dr. Hemang Baxi  
(M) +91-98250 30111

#### Cardiac Surgeons

Dr. Dhiren Shah  
(M)+91-98255 75933  
Dr. Dhaval Naik  
(M)+91-90991 11133  
Dr. Dipesh Shah  
(M)+91-90990 27945

#### Pediatric & Structural

**Heart Surgeons**  
Dr. Shaunak Shah  
(M)+91-98250 44502  
Dr. Ashutosh Singh  
(M)+91-82380 01976

#### Vascular & Endovascular

**Surgeon**  
Dr. Srujal Shah  
(M) +91-91377 88088

#### Cardiac Anaesthetists

Dr. Niren Bhavsar  
(M)+91-98795 71917  
Dr. Hiren Dholakia  
(M)+91-95863 75818  
Dr. Chintan Sheth  
(M)+91-91732 04454

#### Pediatric Cardiologists

Dr. Kashyap Sheth  
(M) +91-99246 12288  
Dr. Milan Chag  
(M) +91-98240 22107

#### Neonatologist and

**Pediatric Intensivist**  
Dr. Amit Chitaliya  
(M)+91-90999 87400

#### Cardiac Electrophysiologist

Dr. Ajay Naik  
(M) +91-98250 82666

#### From the desk of Editor:

Systolic Blood Pressure (SBP) rises progressively with age, while after about the age of 65 there is a modest decline. Isolated Systolic Hypertension (ISH) with increased pulse pressure (Figure 1) is the commonest form of hypertension in the elderly and is a major risk factor for cardiovascular (CV) disease. ISH is defined as SBP >140 mmHg and Diastolic Blood Pressure (DBP) <90 mmHg. It should probably be distinguished from essential hypertension (where SBP and DBP are both increased). Pathogenesis of the two conditions overlaps, but is not identical. There are subtle differences in the approach to treatment. ISH and essential hypertension are not entirely distinct – a patient with essential hypertension may later present with ISH as the DBP decreases with age. However, over 60% of patients with ISH have no preceding hypertension. ISH affects two-thirds of patients over 65 years and three-quarters of patients over 75. It is by far the most common form of hypertension in later life (Figure 2). With the ageing population, this is clearly a very major public health problem.



**Dr. Hemang Baxi**

## Isolated Systolic Hypertension

### Case History

**A 75-year-old man who is a care taker for his disabled wife.**

He is quite active and copes well with household chores. He has high cholesterol and takes a statin as well as aspirin. Blood pressure (BP) has been high for some time, and at the visit is 168/82. The time has come to start treatment, and you wish to discuss this with him. Renal function and creatinine are both normal. Urine is negative on stick testing but he has microalbuminuria on laboratory testing.

### ■ What is the optimal approach to management?

The relative increase in SBP, compared with DBP, relates to decreased vascular compliance with ageing. The latter arises from atherosclerosis, collagen cross-linking and glycosylation, vascular calcification and elastin fragmentation. The aorta and larger vessels lose the ability to expand to accommodate the pulse wave during systole and there is decreased recoil during diastole. Decreased baroreceptor responses and increased salt sensitivity contribute to hypertension in the elderly. Although plasma catecholamines are relatively increased, sensitivity of adrenoreceptors is decreased. Increased

- Is ISH common, and is it important?
- Why does Systolic BP (SBP) selectively increase in the elderly?






Figure-1 : Change in blood pressure with age.




Figure-2 : Presentation of untreated hypertension. IDH, isolated diastolic hypertension; ISH, isolated systolic hypertension; S-D, systolic-diastolic (essential) hypertension. Adapted from Chobanian.

sympathetic drive is, therefore, a less important mechanism and because of this as well as often decreased myocardial function, increased cardiac output is less important than in younger subjects. Activation of the renal artery stenosis is also relatively less important in ISH, explaining why angiotensin-converting enzyme inhibitors and angiotensin receptor blockers are often not the most effective drugs. The addition of angiotensin-converting enzyme inhibitors and angiotensin receptor blockers to treatment is logical where ISH is placing the patient in

danger of end-organ damage. There is particularly strong evidence that these agents may bring about regression of left ventricular hypertrophy and retard the development of nephropathy. Salt sensitivity is increased in older subjects, and this contributes to increased arterial stiffness. Thus, lower salt intake and diuretic therapy are central in management. ISH and essential hypertension are compared in Table 1.

Table-1 : ISH and essential hypertension

	Essential	ISH
Increase cardiac output	+	—
Arterial compliance	+	—
Peripheral resistance	++	+
Left ventricular mass	+	++
Baroreceptor sensitivity	--	—
Plasma catecholamines	+	++
Plasma renin	—	—
Salt sensitivity	+	++

Adapted from Pannarale.

Even a modest decrease in SBP of <5 mmHg reduces cardiac mortality by 7% and stroke mortality by 10%. A target BP of 140/90 seems reasonable for the elderly, but is often not achievable, let alone the tighter target of 130/80, which is recommended for those at higher risk (e.g. those with diabetes and chronic kidney disease). Effective treatment of ISH decreases the risk of CV events by 23%. The landmark trials in this area (see Duprez for a review) were:

- SHEP (1991) followed 4736 patients with ISH for 4.5 years. In treated patients, the rate of non-fatal stroke was decreased by 36%, cardiac disease by 25%, and heart failure by 53%. A more recent (2005) extension of this study following patients for up to 14.3 years showed continuing benefit in terms of event rate reduction. Benefit extended to those with pre-existing diabetes, while those who developed diabetes during the follow-up period but had their BP treated had no



different risk to patients without diabetes. This study was based on the use of chlorthalidone with other agents added as needed.

- Syst-Eur (1997) and Syst-China (1998) followed 4695 and 2394 patients respectively for 2-3 years, with initial therapy based on Calcium Channel Blockers (CCB). These studies showed a reduction in stroke of about 40% and of total CV outcomes of 31-39% for treated patients.

More recent studies have included: LIFE (2002) in which losartan decreased CV out-comes compared with an atenolol-based regimen in 1326 patients with ISH and left ventricular hypertrophy followed for 4.7 years; the Systolic Hypertension in the Elderly Long-term Lacidipine study (SHELL, 2003) in which 1882 patients with ISH were followed for 32 months, showed a 9.3% reduction in event rate with the CCB; the Intervention as a Goal in Hypertension Treatment study (INSIGHT, 2004) which included 1498 patients with ISH followed for 3 years, and showed a 6% decrease in the CV event rate. These, and a number of other short-term studies have, in recent years, highlighted ISH as a clinical problem in the elderly. They have also shown that decreasing BP leads to lower CV risk. Although the various classes of agent are similar in BP-decreasing potency, rennin-angiotensin system blocking drugs and CCBs have the most trial evidence. The former may be slightly superior in preventing stroke and are slightly better tolerated because of the relatively high incidence of peripheral oedema seen with CCBs.

## Treatment of ISH

As with all patients with hypertension, the elderly with ISH should have an overall assessment of CV risk, secondary causes of hypertension should be considered (particularly renovascular) and they should be screened for end-organ damage (eyes, heart and kidneys). Consider also conditions that increase cardiac output and may selectively increase SBP. These include anaemia,

thyrotoxicosis, Paget's disease and aortic regurgitation. Lifestyle modifications should be instituted where possible – maintain or decrease body weight, regular exercise, balanced diet, lower salt intake and avoid excess alcohol. It is assumed that the major advantage of drug treatment is through lowering BP, and none of the major classes has a specific action in ISH. In the elderly, it is important to start with low doses of drugs where possible, and to titrate gradually with careful monitoring of BP response, renal function and electrolytes. Over-vigorous reduction in DBP should be avoided as this may decrease myocardial perfusion.

The first choice of treatment for ISH is low-dose thiazide. The greatest evidence is with hydrochlorthiazide and chlorthalidone. Bendrofluazide is also widely used. The latter is more potent on a milligram for milligram basis and also has a longer half-life (48-72h vs 16-24 h). The disadvantages of thiazides are hypokalemia, increased uric acid, dyslipidaemia, hyperglycaemia, and erectile dysfunction in men. All of these are common in the elderly but less likely to occur with low doses – 12.5 mg of hydrochlorthiazide is a suitable starting dose. Long-acting CCBs should also be considered early in the treatment. These are of proven efficacy and have beneficial effects on vascular remodelling. B-blockers are no longer considered first-line for the elderly patient with ISH, but should certainly be used in those with angina or previous myocardial infarction, and considered in those with heart failure (Figure 3).

## Recent Developments

1. High SBP is very variable in older subjects with reduced vascular compliance. There is an argument for confirming ISH on at least three occasions before the diagnostic label is assigned. A recent large Portuguese study confirmed that ISH was common, particularly after the age of 70. It was not, however, particularly associated with premature mortality, and CV complications often developed very late in life.





2. In a follow-up of participants in the SHEP trial at 14.3 years, a chlorthalidone-based antihypertensive regimen decreased CV mortality – RR 0.86 (95% CI:0.76 to 0.98). Patients who had sustained stroke experienced a particularly poor mortality experience. We should not be over-pessimistic about the dangers of ISH, or over-optimistic about the benefits of treatment. For population-based risk reduction, many patients would have to be treated for many years to appreciably impact on mortality.
3. Over-aggressive treatment of systolic hypertension may lead to an unwanted decrease in DBP. In the elderly, DBP <60 mmHg has been associated with poor prognosis independent of large artery stiffness and left ventricular function. Elderly people taking antihypertensives should be carefully monitored. DBP decreases with age and may warrant altering treatment for ISH.
4. Long-acting CCBs are a very useful adjunct to thiazides as first-line treatment. A recent trial compared amlodipine with a newer CCB manidipine. Chlorthalidone was added where needed. Both CCBs effectively decreased BP. Manidipine had the advantage of a lower incidence of peripheral oedema.
5. The importance of exercise as an intervention for the elderly is often forgotten. Apart from general well-being, exercise has tangible benefits including reducing SBP. Given the now proven benefits of exercise in elderly patients with chronic disease, including diabetes and pulmonary disease, we need to examine how to actively engage elderly people in exercise programmes.

## Conclusions

The vast majority of us will develop hypertension as we age, and ISH is by far the commonest form in older people. Subtly, the pathogenesis is different to that of essential hypertension, which is a disease of younger people. Decreased vascular compliance is the hallmark. The underlying cause of hypertension shifts from more humoral mechanisms in younger subjects to more




Figure-3 : Management of ISH. CCF, congestive cardiac failure; CKD, chronic kidney disease.

mechanical causes in the elderly. This explains why some drugs are less effective in ISH. The condition is important, being strongly linked with adverse CV outcomes, particularly stroke. Treatment with two or more drugs is often needed. Over-vigorous treatment may control ISH but at the expense of undue lowering of DBP, which may reduce coronary perfusion. Low-dose thiazides and long-acting CCBs are the cornerstone of treatment. These should be initiated cautiously and titrated gradually, especially in very aged or frail patients.



## CIMS Clinic Maninagar



### Special Cardiac Check-up rate

<del>Original Package</del>		Discounted Package	
Consultation	₹ 600	Consultation	} Only ₹ 600/-*
Echo	: ₹ 1200	Echo	
ECG	: ₹ 150	ECG	
	<del>: ₹ 1950</del>		

\*This rate valid till July 31 2012 at CIMS Clinic Maninagar only.



### CIMS Clinic (Maninagar )


1<sup>st</sup> Floor, Shant Prabha Heights, Opp. Vallabh Vadi, Bhairavnath Road,  
Maninagar, Ahmedabad-380 008.

For appointment call : +91-79-2544 0381-83 (3 lines)



## CIMS ENT launches Balloon Sinuplasty

A novel & minimal invasive  
treatment for chronic rhinosinusitis



*Balloon Sinuplasty (BSP) is a breakthrough sinus procedure performed to open sinus passages to relieve sinus pain and sinus pressure associated with chronic sinusitis and recurring sinus infections.*

#### Contact for appointment

Dr. Navin K. Patel (M.S. ENT)  
Mobile : +91-98240 23746

Dr. Monark B. Shah (M.S. ENT)  
Mobile : +91-98244 44338



#### The benefits of Balloon Sinuplasty include:

- **Safe** : Many patients have been treated safely with Balloon Sinuplasty (without open surgery).
- **Proven** : Over 95% of patients who've had the procedure say they would have it again.
- **Fast** : While recovery time varies with each patient, they can return quickly to normal activities.



## Peripheral Vascular Disease (PVD) Workshop by Dr. Ashit Jain

**August 31 - September 1, 2012**

**Patients who are eligible :** Carotid Artery Stenosis • Renal Artery Stenosis • Acute Limb Ischemia • Critical Limb Ischemia • Claudication • Aortoiliac occlusive disease • Femoropopliteal Disease • Brachiocephalic Arterial Disease • Venous Thromboembolic Disease • Thoracic Abdominal Aortic Aneurysms • Mesenteric Disease • Catheter-Based Interventions for Failing Hemodialysis Accesses • Infrapopliteal Peripheral Arterial Disease • Intracranial Arterial Stenotic Disease • Vertebral Arterial Disease

**Patients will be provided following FREE services:**

**1. Consultation 2. ABI**

**Daily screening camp of the concerned patients will be held in the month of July, 2012 at CIMS Hospital. Time : 2.00 pm - 6.00 pm**

Organized by  
**CIMS**  
Care Institute of Medical Sciences  
At CIMS... we care



Dr. Ashit Jain is a well known Interventional Cardiologist practicing for the past 20 years in California, USA. Graduated from University of Delhi, completed Fellowship in Interventional Cardiology and Peripheral Vascular Disease at Ochsner Medical Center in New Orleans, USA, he has developed an extensive clinical research program at Washington Hospital in Fremont, California and is involved in multiple new device research technologies. He has also served as site principal investigator on over 26 multi-center clinical research trials and has written and presented many abstracts and publications in the field. A pioneer in Carotid Interventional Programs in the San Francisco Bay area, he is affiliated with five hospitals in the East Bay of San Francisco and has personally performed over 500 carotid interventions.

**You may call any of our cardiologists listed on the front page to facilitate**

## ECP (External Counter Pulsation Therapy)



**No surgery. No intervention. No pain.**

### Advantages of ECP therapy over other options:

- A non-invasive treatment for coronary artery stenosis.
- Performed on an out patient basis.
- Safe, painless, comfortable with no side effects.
- Cost effective.



**Non-invasive alternative to treat heart disease\***

\*for patients with heart blocks in arteries who cannot or don't want to undergo angioplasty / bypass & for old bypass patients having recurrence of chest pain



- ECP (External Counter Pulsation), is a non-invasive procedure which reduces symptoms of angina pectoris.
- ECP therapy is clinically tested and proven atraumatic outpatient alternative procedure to standard surgical interventions, namely balloon angioplasty (PTCA) and bypass surgery (CABG).

**You may call any of our cardiologists listed on the front page to facilitate**

**CIMS**<sup>®</sup>  
Care Institute of Medical Sciences  
At CIMS... we care

For appointment call : +91-79-3010 1200, 3010 1008  
Mobile : +91-98250 66661 or email on [opd.rec@cims.me](mailto:opd.rec@cims.me)





## CIMS-CON

### 2013

Education For Innovation

January 4-6, 2013

Organized by

**CIMS**  
Care Institute of Medical Sciences



**The Heart Care Clinic**  
Clinical Care Consultants

9<sup>th</sup> Annual Scientific Symposium, 18<sup>th</sup> Year of Academics

**SUPER  
EARLY BIRD  
REGISTRATION**

**₹ 2,500/- only\***  
**instead of ₹ 5,000/-**

**The first 500 registered delegates  
will get a FREE DVD Set of  
CIMS-CON 2012**

**And also 5 more lucky draw winners will be gifted iPad / TV / DVD Player**

Only DVD set without CIMS-CON 2013 registration is available for ₹ 2,000/-

\* Offer valid till May 31, 2012

### CIMS-CON 2013 Registration Form

Cheque or DD's to be made A/C payee and in the name of 'CIMS Hospital Pvt. Ltd.' Kindly mail the registration form along with the cheque/DD to our office. All Cash Payment are to be made at CIMS Hospital, Ahmedabad only.

**Please note that it is mandatory to provide all the information. Please fill all fields in CAPITAL LETTERS**

☐ Prof. ☐ Dr. ☐ Mr. ☐ Mrs. ☐ Ms.

Full Name

Qualification

Resi. Address

City  Pin Code

Phone (STD code)  Mobile

Email

\*\* Hotel Accommodation is optional. If you have applied for accommodation, please send a separate deposit cheque of ₹ 3000 to cover the cost of your stay for two nights. Spouse hotel registration will be charged extra. Students also need to pay for Hotel Accommodation at the same rate.

**Hotel accommodation will be allotted on first come, first served basis as per preference of doctors from our list of hotels.**

#### Payment Details

₹

₹ in word :

DD/Cheque No.

Date

Bank :

#### Payment Details - Spouse Hotel Registration\*

**Spouse Hotel Registration Fee ₹ 3,500/- (Non Refundable)**

Delegate Name :

Spouse Name :

Contact No. :

₹

₹ in word :

DD/Cheque No.

Date

Bank :

\*Please give separate cheque of ₹ 3,500/- for spouse hotel registration

**CIMS**  
Care Institute of Medical Sciences

CIMS Hospital, Nr. Shukan Mall, Off Science City Road, Sola, Ahmedabad-380060

Phone : +91-79-3010 1059 / 1060 Fax: +91-79-2771 2770 (M) +91-98250 66664, 98250 66668

Email : [cimscon@cims.me](mailto:cimscon@cims.me), [www.cimscon.com](http://www.cimscon.com) / [www.cims.me](http://www.cims.me)



Healthy Heart Registered under **RNI No. GUJENG/2008/28043**

Permitted to post at PSO, Ahmedabad-380002 on the 12<sup>th</sup> to 17<sup>th</sup> of every month under  
Postal Registration No. **GAMC-1725/2012-2014** issued by SSP Ahmedabad valid upto 31<sup>st</sup> December, 2014  
Licence to Post Without Prepayment No. **CPMG/GJ/97/2012** valid upto 30<sup>th</sup> June, 2012

**If undelivered Please Return to :**

CIMS Hospital, Nr. Shukan Mall,  
Off Science City Road, Sola, Ahmedabad-380060.  
Ph. : +91-79-2771 2771-75 (5 lines)  
Fax: +91-79-2771 2770  
Mobile : +91-98250 66664, 98250 66668

**Subscribe "Healthy Heart"** : Get your "Healthy Heart", the information of the latest medical updates only ₹ 60/- for one year.  
To subscribe pay ₹ 60/- in cash or cheque/DD at CIMS Hospital Pvt. Ltd. Nr. Shukan Mall, Off Science City Road, Sola, Ahmedabad-380060. Phone : +91-79-3010 1059 / 3010 1060. Cheque/DD should be in the name of : **"CIMS Hospital Pvt. Ltd."**  
Please provide your **complete postal address with pincode, phone, mobile and email id** along with your subscription

## CARE INSTITUTE OF MEDICAL SCIENCES



A premier multi-super specialty  
**GREEN Hospital**



CIMS Hospital : Nr. Shukan Mall, Off Science City Road, Sola, Ahmedabad-380060. Ph.:  
+91-79-2771 2771-75 (5 lines) email : [info@cims.me](mailto:info@cims.me) web : [www.cims.me](http://www.cims.me)  
For appointment call : +91-79-3010 1200, 3010 1008  
Mobile : +91-98250 66661 or email on [opd.rec@cims.me](mailto:opd.rec@cims.me)

**Ambulance & Emergency : +91-98244 50000, 97234 50000, 90990 11234**

Printed, Published and Edited by Dr. Keyur Parikh on behalf of the CIMS Hospital  
Printed at Hari Om Printery, 15/1, Nagori Estate, Opp. E.S.I. Dispensary, Dudheshwar Road, Ahmedabad-380004.  
Published from CIMS Hospital, Nr. Shukan Mall, Off Science City Road, Sola, Ahmedabad-380060.

