



HEALTHY HEART

VOLUME-10 | ISSUE-120 | NOVEMBER 05, 2019

Honorary Editor :
Dr. Hemang Baxi



Systolic Blood Pressure (SBP) rises progressively with age, while after about the age of 65 there is a modest decline. Isolated Systolic Hypertension (ISH) with increased pulse pressure (Figure 1) is the commonest form of hypertension in the elderly and is a major risk factor for cardiovascular (CV) disease. ISH is defined as SBP >140 mmHg and Diastolic Blood Pressure (DBP) <90 mmHg. It should probably be distinguished from essential hypertension (where SBP and DBP are both increased). Pathogenesis of the two conditions overlaps, but is not identical. There are subtle differences in the approach to treatment. ISH and essential hypertension are not entirely distinct – a patient with essential hypertension may later present with ISH as the DBP decreases with age. However, over 60% of patients with ISH have no preceding hypertension. ISH affects two-thirds of patients over 65 years and three-quarters of patients over 75. It is by far the most common form of hypertension in later life (Figure 2). With the ageing population, this is clearly a very major public health problem.

Isolated Systolic Hypertension

Case History

A 75-year-old man who is a care taker for his disabled wife.

He is quite active and copes well with household chores. He has high cholesterol and takes a statin as well as aspirin. Blood pressure (BP) has been high for some time, and at the visit is 168/82. The time has come to start treatment, and you wish to discuss this with him. Renal function and creatinine are both normal. Urine is negative on stick testing but he has microalbuminuria on laboratory testing.

- Is ISH common, and is it important?
- Why does Systolic BP (SBP) selectively increase in the elderly?
- What is the optimal approach to management?

The relative increase in SBP, compared with DBP, relates to decreased vascular compliance with ageing. The latter arises from atherosclerosis, collagen cross-linking and glycosylation, vascular calcification and elastin fragmentation. The aorta and larger

vessels lose the ability to expand to accommodate the pulse wave during systole and there is decreased recoil during diastole. Decreased baroreceptor responses and increased salt sensitivity contribute to hypertension in the elderly. Although plasma catecholamines are relatively increased, sensitivity of adrenoceptors is decreased. Increased




Figure-1 : Change in blood pressure with age.

sympathetic drive is, therefore, a less important mechanism and because of this as well as often decreased myocardial function, increased cardiac output is less important than in younger subjects. Activation of the

Cardiologists

Dr. Satya Gupta (M) +91-99250 45780	Dr. Milan Chag (M) +91-98240 22107
Dr. Vineet Sankhla (M) +91-99250 15056	Dr. Urmil Shah (M) +91-98250 66939
Dr. Vipul Kapoor (M) +91-98240 99848	Dr. Hemang Baxi (M) +91-98250 30111
Dr. Tejas V. Patel (M) +91-89403 05130	Dr. Anish Chandarana (M) +91-98250 96922
Dr. Gunvant Patel (M) +91-98240 61266	Dr. Ajay Naik (M) +91-98250 82666
Dr. Keyur Parikh (M) +91-98250 26999	

Congenital & Structural Heart Disease Specialist

Dr. Kashyap Sheth (M) +91-99246 12288	Dr. Milan Chag (M) +91-98240 22107
Dr. Divyesh Sadadiwala (M) +91-8238339980	

Cardiothoracic & Vascular Surgeons

Dr. Dhiren Shah (M) +91-98255 75933
Dr. Dhaval Naik (M) +91-90991 11133
Dr. Amit Chandan (M) +91-96990 84097

Paediatric & Structural Heart Surgeons

Dr. Shaunak Shah (M) +91-98250 44502

Cardiovascular, Thoracic & Thoracoscopic Surgeon

Dr. Pranav Modi (M) +91-99240 84700

Cardiac Anaesthetists

Dr. Niren Bhavsar (M) +91-98795 71917
Dr. Hiren Dholakia (M) +91-95863 75818
Dr. Chintan Sheth (M) +91-91732 04454

Cardiac Electrophysiologist

Dr. Ajay Naik (M) +91-98250 82666
Dr. Vineet Sankhla (M) +91-99250 15056

Neonatologist and Paediatric Intensivist

Dr. Amit Chitaliya (M) +91-90999 87400
--



renal artery stenosis is also relatively less important in ISH, explaining why angiotensin-converting enzyme inhibitors and angiotensin receptor blockers are often not the most effective drugs. The addition of angiotensin-converting enzyme inhibitors and angiotensin receptor blockers to treatment is logical where ISH is placing the patient in




Figure-2 : Presentation of untreated hypertension. IDH, isolated diastolic hypertension; ISH, isolated systolic hypertension; S-D, systolic-diastolic (essential) hypertension. Adapted from Chobanian.

danger of end-organ damage. There is particularly strong evidence that these agents may bring about regression of left ventricular hypertrophy and retard the development of nephropathy. Salt sensitivity is increased in older subjects, and this contributes to increased arterial stiffness. Thus, lower salt intake and diuretic therapy are central in management. ISH and essential hypertension are compared in Table 1.

Even a modest decrease in SBP of <5 mmHg reduces cardiac mortality by 7%

Table-1 : ISH and essential hypertension

	Essential	ISH
Increase cardiac output	+	—
Arterial compliance	+	—
Peripheral resistance	++	+
Left ventricular mass	+	++
Baroreceptor sensitivity	—	—
Plasma catecholamines	+	++
Plasma renin	—	—
Salt sensitivity	+	++
Adapted from Pannarale.		

and stroke mortality by 10%. A target BP of 140/90 seems reasonable for the elderly, but is often not achievable, let alone the tighter target of 130/80, which is recommended for those at higher risk (e.g. those with diabetes and chronic kidney disease). Effective treatment of ISH decreases the risk of CV events by 23%. The landmark trials in this area (see Duprez for a review) were:

- SHEP (1991) followed 4736 patients with ISH for 4.5 years. In treated patients, the rate of non-fatal stroke was decreased by 36%, cardiac disease by 25%, and heart failure by 53%. A more recent (2005) extension of this study following patients for up to 14.3 years showed continuing benefit in terms of event rate reduction. Benefit extended to those with pre-existing diabetes, while those who

developed diabetes during the follow-up period but had their BP treated had no different risk to patients without diabetes. This study was based on the use of chlorthalidone with other agents added as needed.

- Syst-Eur (1997) and Syst-China (1998) followed 4695 and 2394 patients respectively for 2-3 years, with initial therapy based on Calcium Channel Blockers (CCB). These studies showed a reduction in stroke of about 40% and of total CV outcomes of 31-39% for treated patients.

More recent studies have included: LIFE (2002) in which losartan decreased CV out-comes compared with an atenolol-based regimen in 1326 patients with ISH and left



ventricular hypertrophy followed for 4.7 years; the Systolic Hypertension in the Elderly Long-term Lacidipine study (SHELL, 2003) in which 1882 patients with ISH were followed for 32 months, showed a 9.3% reduction in event rate with the CCB; the Intervention as a Goal in Hypertension Treatment study (INSIGHT, 2004) which included 1498 patients with ISH followed for 3 years, and showed a 6% decrease in the CV event rate. These, and a number of other short-term studies have, in recent years, highlighted ISH as a clinical problem in the elderly. They have also shown that decreasing BP leads to lower CV risk. Although the various classes of agent are similar in BP-decreasing potency, rennin-angiotensin system blocking drugs and CCBs have the most trial evidence. The former may be slightly superior in preventing stroke and are slightly better tolerated because of the relatively high incidence of peripheral oedema seen with CCBs.

Treatment of ISH

As with all patients with hypertension, the elderly with ISH should have an overall assessment of CV risk, secondary causes of hypertension should be considered (particularly renovascular) and they should be screened for end-organ damage (eyes, heart and kidneys). Consider also conditions that increase cardiac output and may selectively increase SBP. These include anaemia,

thyrotoxicosis, Paget's disease and aortic regurgitation. Lifestyle modifications should be instituted where possible – maintain or decrease body weight, regular exercise, balanced diet, lower salt intake and avoid excess alcohol. It is assumed that the major advantage of drug treatment is through lowering BP, and none of the major classes has a specific action in ISH. In the elderly, it is important to start with low doses of drugs where possible, and to titrate gradually with careful monitoring of BP response, renal function and electrolytes. Over-vigorous reduction in DBP should be avoided as this may decrease myocardial perfusion.

The first choice of treatment for ISH is low-dose thiazide. The greatest evidence is with hydrochlorthiazide and chlorthalidone. Bendrofluazide is also widely used. The latter is more potent on a milligram for milligram basis and also has a longer half-life (48-72h vs 16-24 h). The disadvantages of thiazides are hypokalemia, increased uric acid, dyslipidaemia, hyperglycaemia, and erectile dysfunction in men. All of these are common in the elderly but less likely to occur with low doses – 12.5 mg of hydrochlorthiazide is a suitable starting dose. Long-acting CCBs should also be considered early in the treatment. These are of proven efficacy and have beneficial effects on vascular remodelling. B-blockers are no longer considered first-line for the elderly patient with ISH, but should

certainly be used in those with angina or previous myocardial infarction, and considered in those with heart failure (Figure 3).

Recent Developments

1. High SBP is very variable in older subjects with reduced vascular compliance. There is an argument for confirming ISH on at least three occasions before the diagnostic label is assigned. A recent large Portuguese study confirmed that ISH was common, particularly after the age of 70. It was not, however, particularly associated with premature mortality, and CV complications often developed very late in life.
2. In a follow-up of participants in the SHEP trial at 14.3 years, a chlorthalidone-based antihypertensive regimen decreased CV mortality – RR 0.86 (95% CI: 0.76 to 0.98). Patients who had sustained stroke experienced a particularly poor mortality experience. We should not be over-pessimistic about the dangers of ISH, or over-optimistic about the benefits of treatment. For population-based risk reduction, many patients would have to be treated for many years to appreciably impact on mortality.
3. Over-aggressive treatment of systolic hypertension may lead to an unwanted decrease in DBP. In the elderly, DBP <60 mmHg has been associated with poor



prognosis independent of large artery stiffness and left ventricular function. Elderly people taking antihypertensives should be carefully monitored. DBP decreases with age and may warrant altering treatment for ISH.

4. Long-acting CCBs are a very useful adjunct to thiazides as first-line treatment. A recent trial compared amlodipine with a newer CCB manidipine. Chlorthalidone was added where needed. Both CCBs effectively decreased BP. Manidipine had the advantage of a lower incidence of peripheral oedema.
5. The importance of exercise as an intervention for the elderly is often forgotten. Apart from general well-being, exercise has tangible benefits including reducing SBP. Given the now proven benefits of exercise in elderly patients with chronic disease, including diabetes and pulmonary disease, we need to examine how to actively engage elderly people in exercise programmes.

younger subjects to more mechanical causes in the elderly. This explains why some drugs are less effective in ISH. The condition is important, being strongly linked with adverse CV outcomes, particularly stroke. Treatment with two or more drugs is often needed. Over-vigorous

treatment may control ISH but at the expense of undue lowering of DBP, which may reduce coronary perfusion. Low-dose thiazides and long-acting CCBs are the cornerstone of treatment. These should be initiated cautiously and titrated gradually, especially in very aged or frail patients.




Figure-3 : Management of ISH. CCF, congestive cardiac failure; CKD, chronic kidney disease.

Conclusions

The vast majority of us will develop hypertension as we age, and ISH is by far the commonest form in older people. Subtly, the pathogenesis is different to that of essential hypertension, which is a disease of younger people. Decreased vascular compliance is the hallmark. The underlying cause of hypertension shifts from more humoral mechanisms in



CIMS HOSPITAL


GASTROENTEROLOGY

CENTRE OF EXCELLENCE - GASTROENTEROLOGY FACILITIES

STATE-OF-THE-ART - PENTAX
FIRST IN GUJARAT HD EUS & EBUS (J 10 SERIES) WITH HD OPTIVISTA
PROCESSOR FOR BETTER DIAGNOSIS.



Upper GI Endoscopy



Lower GI Endoscopy



ERCP




Endoscopic Ultrasound




Fibroscan




Capsule Endoscopy



Esophageal - Manometry



Anorectal Manometry



24 hr pH Study



Colon Transit Test



MRI Defecography



Breath Test

GASTROENTEROLOGY TEAM

Dr. BHAVESH THAKKAR
(Gastroenterologist)
M : +91 97277 07214

DR. ABHINAV JAIN
(Gastroenterologist)
M : +91 7666373288





HEALTHY HEART

VOLUME-10 | ISSUE-120 | NOVEMBER 05, 2019

FOR THE FIRST TIME WOMEN SESSION AT JIC (JANUARY 3-4, 2020)

TO ACKNOWLEDGE THE GROWING
CONTRIBUTION OF WOMEN
TO THE MEDICAL FIELD

**REGISTER
TODAY**
FOR A SPECIAL
SURPRISE GIFT
LIMITED SEATS
AVAILABLE

ATTEND THE SESSION

NETWORKING

ADVANCED LEARNING & INTERACTIVE SESSION

LIVE CASE BASED PRESENTATIONS

SPECIAL PRIZES ELIGIBILITY CRITERIA

- Women MD (Physician) with MD (Medicine) degree
- Working as MD Physicians full time or part time for 1 year
- Consulting / Seeing at least 20 patients a month
- Committed to attend all conference days
- PG MD (Medicine) women students will also be accepted.

CIMS Hospital, Off. Science City Road, Sola, Ahmedabad - 380060.

Ph.: +91-79-2772 1059/1060

Email : communication@cimshospital.org | www.jicindia.org

YOU WILL RECEIVE

Free certificate
to attend 1
CIMS Learning Centre
(CLC)
Course in
2020

Surprise
Felicitation Gift

25% discount
for
ECHO Fellowship
Course

Organized by

GMERS
Medical College,
Sola,
Ahmedabad

Supported by



CIMSRE

Care Institute
Medical Society
for Research
and Education

In Association with



Conference Secretariat

CIMS Hospital, Off. Science City Road, Sola, Ahmedabad - 380060.

Ph.: +91-79-2772 1059/1060

Fax : +91-79-2771 2770 (M) +91-98250 66664, 98250 66668

Email : communication@cimshospital.org | www.jicindia.org



Highest Lectures Ever.
Discussion. Debates. Interaction

JIC 2020

Joint International Conference

January 3-5, 2020

Register Today for
Entire 3-day JIC @ Rs. 6500/- Only
(upto December 31, 2019)

visit www.jicindia.org
for more information



**CIMS
Learning
Centre
2020**

For any query, please email on :
clc@cimshospital.org

Echocardiography Fellowship Programme

Registration Fees : ₹ 35,000/-*

Course Directors : Dr. Satya Gupta / Dr. Milan Chag / Dr. Tejas V. Patel / Dr. Vineet Sankhla
Date : One Week (Monday to Saturday)
Duration : 6 days
Number of Seats : 15
Venue : CIMS Auditorium

Programme Overview:

Sights and Sounds of Echocardiography at CIMS Hospital is a premier education course for Echocardiologists, Fellows in training, and Physicians in clinical practice. Over the past few years, we have trained and taught more than 1000 candidates. This unique programme is intended for those who want to learn Echocardiography right from basics. Overall programme includes lectures on basics of Echocardiography, theoretical and practical aspects of Ischemic, Valvular and Congenital Heart Disease in relation to Echocardiography and hands-on supervised training. This is the star educational programme of the CIMS Learning Centre and has been appreciated worldwide.

Online registration & payment on
[www.cims.org /clc](http://www.cims.org/clc)

Programme Highlights:

- 6 days extensive teaching of echocardiography
- Easy access to faculty members
- Supervised hands-on practice
- Provision of course materials (Textbook of Echocardiography)

Schedule of Echocardiography Fellowship 2020

Month	Start Date	End Date
February	24-02-2020	29-02-2020
April	20-04-2020	25-04-2020
June	22-06-2020	27-06-2020
September	24-08-2020	29-08-2020
November	19-10-2020	24-10-2020

> Certificate of attendance will be given at the end of the course.

* Hotel Registration Separate



HEALTHY HEART

VOLUME-10 | ISSUE-120 | NOVEMBER 05, 2019

Healthy Heart Registered under RNI No. GUJENG/2008/28043

Published on 5th of every month

Permitted to post at PSO, Ahmedabad-380002 on the 12th to 17th of every month under
Postal Registration No. **GAMC-1725/2018-2020** issued by SSP Ahmedabad valid upto 31st December, 2020
Licence to Post Without Prepayment No. **PMG/HQ/055/2018-20** valid upto 31st December, 2020

If undelivered Please Return to :

CIMS Hospital, Nr. Shukan Mall,
Off Science City Road, Sola, Ahmedabad-380060.
Ph. : +91-79-2771 2771-72
Fax: +91-79-2771 2770
Mobile : +91-98250 66664, 98250 66668

Subscribe "Healthy Heart" : Get your "Healthy Heart", the information of the latest medical updates only ` 60/- for one year.
To subscribe pay ` 60/- in cash or cheque/DD at CIMS Hospital Pvt. Ltd. Nr. Shukan Mall, Off Science City Road, Sola,
Ahmedabad-380060. Phone : +91-79-2772 1059 / 2772 1060. Cheque/DD should be in the name of : "CIMS Hospital Pvt. Ltd."
Please provide your complete postal address with pincode, phone, mobile and email id along with your subscription



CIMS Learning Centre

Skills Development Centre

Advanced ECG Learning Course

November 24, 2019 (Sunday)

Course Directors : Dr. Ajay Naik / Dr. Vipul Kapoor / Dr. Tejas V. Patel / Dr. Vineet Sankhla
Duration : 1 day
Number of Seats : 50
Venue : CIMS Auditorium

Programme Overview:

Advanced ECG course is designed for Practicing Physicians having basic knowledge of ECG. This programme covers a wide range of ECG lectures and tips to identify various type of simple & complex Arrhythmias and rare ECG abnormalities.

The aim of this course is to present complex various ECG diagnosis pattern in a simple and easy-to-understand manner.

Online registration & payment on www.cims.org/clc

Registration Fees: ` 500/- | Spot Registration Fees: ` 1,000/-

Non-refundable

For any query, please email on : clc@cimshospital.org

> Certificate of attendance will be given at the end of the course.

CIMS Hospital : Regd Office: Plot No.67/1, Opp. Panchamrut Bungalows, Nr. Shukan Mall, Off Science City Road, Sola, Ahmedabad - 380060.

Ph. : +91-79-2771 2771-72 Fax: +91-79-2771 2770.

CIMS Hospital Pvt. Ltd. | CIN : U85110GJ2001PTC039962 | info@cims.org | www.cims.org

Printed, Published and Edited by Dr. Keyur Parikh on behalf of the CIMS Hospital

Printed at Hari Om Printery, 15/1, Nagori Estate, Opp. E.S.I. Dispensary, Dudheshwar Road, Ahmedabad-380004.

Published from CIMS Hospital, Nr. Shukan Mall, Off Science City Road, Sola, Ahmedabad-380060.